

Published in final edited form as:

J Am Chem Soc. 2007 February 28; 129(8): 2216–2217. doi:10.1021/ja0682576.

Fluorescent Sequence-Specific dsDNA Binding Oligomers

David M. Chenoweth, Anne Viger, and Peter B. Dervan*

Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125

Sequence-specific detection methods for double-stranded DNA (dsDNA) that obviate the need for denaturation would provide useful tools for bioorganic chemistry and genetics.¹ Previous efforts, such as molecular beacons² or peptide nucleic acid–thiazole orange (PNA–TO) conjugates,³ require harsh denaturation conditions for hybridization to single-stranded DNA.^{2–4} Previous efforts from our laboratory for the sequence-specific detection of dsDNA have focused on pyrrole–imidazole (Py–Im) polyamide–fluorophore conjugates, such as tetramethylrhodamine (TMR) or thiazole orange (TO), that bind in the minor groove of DNA.^{5–7} TMR fluorescence was shown to be quenched when the fluorophore was covalently linked to the ring nitrogen of a pyrrole recognition element within a polyamide.⁵ Remarkably, fluorescence was restored in a sequence-dependent manner upon binding to dsDNA.⁵ Similarly, polyamide–TO intercalator conjugates also demonstrate fluorescence enhancement in the presence of match dsDNA.⁶

Having established Py–Im polyamide–dye conjugates as a suitable platform for sequence-specific fluorescent dsDNA detection,^{5,6} we sought to develop a new class of fluorescent DNA binders wherein the fluorescent moiety is an integrated part of the recognition modules. We report here the design of sequence-specific fluorescent dsDNA-binding oligomers (Figure 1) which incorporate multiple 6-5 fused dimer recognition modules⁸ and show a marked fluorescent enhancement upon excitation at 340 nm in the presence of dsDNA. Oligomer **O1** contains the chlorothiophene-benzimidazole (Ct–Bi–), pyrrole-imidazopyridine (–Py–Ip–), and imidazole-hydroxybenzimidazole (–Im–Hz–) recognition modules, whereas oligomer **O2** contains imidazole-imidazopyridine (Im–Ip–) and two pyrrole-benzimidazole (–Py–Bi–) modules. The binding affinities of **O1** and **O2** targeted to two biologically important sequences, 5'-ATACGT-3' (**O1**) and 5'-WGGGGW-3' (**O2**), were determined to be $K_a = 1.6 \times 10^9 \text{ M}^{-1}$ and $2.6 \times 10^9 \text{ M}^{-1}$, respectively, by quantitative DNase I footprinting.^{9–11}

A library of dsDNA hairpins containing six base-pair match and mismatch binding sites for **O1** and **O2** was used to investigate their emission properties (Figure 2). The dsDNA library for **O1** and **O2** contained match sites (**1** and **8**, respectively), single base-pair (bp) mismatch sites (**2–5** and **10–12**, respectively), double bp mismatch sites (**6** and **13**, respectively), and full mismatch sites (**7** for both oligomers). The dsDNA **9** contains the 4-G match site of oligomer **O2**; however, the flanking sequence has been changed to emphasize the effect on binding. The presence of G · C bp under the tail is expected to lower the binding affinity of **O2** as compared to that of dsDNA **8**.¹²

Oligomers **O1** and **O2**, (1 μM concentration) were each incubated with an increasing concentration (1 nM to 1 μM) of dsDNA, and their emission spectra were recorded after

dervan@caltech.edu.

Supporting Information Available: Experimental procedures, and spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

excitation at 340 nm. The oligomers exhibited a marked increase in fluorescence upon addition of dsDNA containing their match site **1** and **8**, respectively (Figures 3 and 4).¹³ Oligomer **O1** showed a moderate decrease in fluorescence intensity in the presence of dsDNA **2**, but proved to be much more sensitive to the incorporation of single base-pair mismatches at the alternate positions in dsDNAs **3–5** (Figure 4a). The incorporation of multiple base-pair mismatches in dsDNAs **6** and **7** showed a significant diminution in fluorescence intensity for **O1**. Oligomer **O2** exhibited a similar trend in sequence specificity, with a moderate decrease in fluorescence intensity observed upon incorporation of single base-pair mismatches (**9–12**) and a more significant decrease with multiple mismatches (**13** and **7**, Figure 4b).

Sequence-specific DNA binding molecules containing a fluorescent switch integrated as part of the recognition modules provides a method to detect DNA sequences without denaturation and in the absence of conjugation to a dye molecule. Fluorescent oligomers may be useful as site-specific chromosome paints for telomeric and centromeric repeats^{14a,b} and could provide insight into cellular trafficking of DNA binding compounds.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Bert T. Lai in the Beckman Institute Laser Resource Center for use of the fluorimeter. We are grateful to The National Institutes of Health for grant support and the Swiss National Science Foundation for a fellowship to A.V.

References

1. Chan EY, Goncalves NM, Haeusler RA, Hatch AJ, Larson JW, Maletta AM, Yantz GR, Carstea ED, Fuchs M, Wong GG, Sullans SR, Gilmanshin R. *Genome Res.* 2004; 14:1137–1146. [PubMed: 15173119]
2. (a) Tyagi S, Kramer FR. *Nat Biotechnol.* 1996; 14:303–308. [PubMed: 9630890] (b) Tyagi S, Bratu DP, Kramer FR. *Nat Biotechnol.* 1998; 16:49–53. [PubMed: 9447593] (c) Kostrikis LG, Tyagi S, Mhlanga MM, Ho DD, Kramer FR. *Science.* 1998; 279:1228–1229. [PubMed: 9508692]
3. Svanvik N, Westman G, Wang D, Kubista M. *Anal Biochem.* 2000; 281:26–35. [PubMed: 10847607]
4. (a) Whitcombe D, Theaker J, Guy SP, Brown T, Little S. *Nat Biotechnol.* 1999; 17:804–807. [PubMed: 10429248] (b) Thelwell N, Millington S, Solinas A, Booth J, Brown T. *Nucleic Acids Res.* 2000; 28:3752–3761. [PubMed: 11000267] (c) Jenkins Y, Barton JK. *J Am Chem Soc.* 1992; 114:8736–8738.
5. (a) Rucker VC, Foister S, Melander C, Dervan PB. *J Am Chem Soc.* 2003; 125:1195–1202. [PubMed: 12553822] (b) Rucker VC, Dunn AR, Sharma S, Dervan PB, Gray HB. *J Phys Chem B.* 2004; 108:7490–7494.
6. Fechter EJ, Olenyuk B, Dervan PB. *J Am Chem Soc.* 2005; 127:16685–16691. [PubMed: 16305259]
7. Approaches based on peptide–thiazole orange conjugates show enhanced fluorescence in the presence of calf thymus (CT) DNA; however, DNA sequence specificity has not been observed. Carreon JR, Mahon KP Jr, Kelley SO. *Org Lett.* 2004; 6:517–519. [PubMed: 14961612]
8. Doss R, Marques MA, Foister S, Chenoweth DM, Dervan PB. *J Am Chem Soc.* 2006; 128:9074–9079. [PubMed: 16834381]
9. Viger A, Dervan PB. *Bioorg Med Chem.* 2006; 14:8539–8549. [PubMed: 16971129]
10. Chenoweth DM, Poposki JA, Marques MA, Dervan PB. *Bioorg Med Chem.* 2007; 15:759–770. [PubMed: 17095230]

11. The sequence 5'-ATACGT-3' is the hypoxia response element in the VEGF promotor, and 5'-WGGGGW-3' is a sequence within the NF- κ B response element.^{9,10}
12. For a study of flanking sequence effects see: Swalley SE, Baird EE, Dervan PB. *J Am Chem Soc.* 1999; 121:1113–1120.
13. Quenching in the absence of DNA could be attributed to intramolecular quenching or solvent-assisted quenching. Studies are underway to elucidate the mechanism.
14. (a) Gygi MP, Ferguson MD, Mefford HC, Lund KP, O'Day C, Zhou P, Friedman C, Engh G, Stolowitz ML, Trask BJ. *Nucleic Acids Res.* 2002; 30:2790–2799. [PubMed: 12087162] (b) Blattes R, Monod C, Susbielle G, Cuvier O, Wu J, Hsieh T, Laemmli UK, Kas E. *EMBO J.* 2006; 25:2397–2408. [PubMed: 16675949]





Figure 1. Structure of oligomers. (a) Oligomer **O1** containing Ct-Bi-, -Py-Ip-, and -Im-Hz- recognition modules. (b) Oligomer **O2** containing Im-Ip- and two -Py-Bi- recognition modules.

**Figure 2.**

Design of dsDNA library. (a) dsDNA sequences used for **O1**. (b) dsDNA sequences used for **O2**.




Figure 3. Fluorescence emission spectra of **O1** and **O2** (1 μM) after 12 h incubation with their match binding site dsDNA (λ_{Ex} = 340 nm). (a) Data for compound **O1**. (b) Data for compound **O2**. The emission was shown to plateau beyond 1 equiv DNA. (See Supporting Information for plots.)




Figure 4. Plot of dsDNA concentration versus normalized fluorescence for each dsDNA. (a) Data for compound **O1**. (b) Data for compound **O2**.